

Asymptotic Behaviour of Liquid-Liquid Dispersions

Christoph Walker

Institut für Mathematik, Universität Zürich-Irchel,
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland
(cwalker@math.unizh.ch)

(MS received 23 Septemeber 2003; 18 February 2004)

Based on earlier results on existence, we study the asymptotic behaviour of solutions to the coalescence-breakage equations including the volume scattering phenomenon and high energy collisions. The solutions are shown to converge towards one particular equilibrium provided the kernels satisfy a kind of reversibility. We also derive stability of these equilibria in a suitable topology.

1. Introduction

In the present article we consider the evolution of a liquid-liquid dispersion, which is a system formed by two immiscible liquids, and where one of these liquids consists of a very large number of droplets that are finely distributed in the other one. These droplets undergo then the influences of binary coalescence and binary breakage meaning that two droplets can merge to build a larger droplet or that a droplet can split into two smaller ones.

Different from most other models considered in literature, we take into account that droplets cannot become arbitrarily large and that experimental observations suggest the existence of a maximal droplet mass (or volume) beyond which no droplet can survive (see [22]). A particular model paying attention to this feature was introduced for the first time by Fasano and Rosso [13] (see also [14], [21], or [4]) and was then developed further by the author [27]. Such a maximal droplet size requires a new interaction mechanism, called *volume scattering*, to prevent the occurrence of droplets that are "too large". The underlying idea is that if two droplets collide having a cumulative mass exceeding the maximal droplet mass, the virtual droplet is highly unstable and immediately decays into two droplets both with mass within the admissible range.

Another new feature taken into consideration in our model is the possibility of high energy collisions leading to a shattering of the involved droplets. Such a breakage mode has been contemplated in physical literature (cf. [7], [8], or [29]) but — at least up to the author's knowledge — only its discrete version has been investigated mathematically so far (see [20]).

We describe the evolution of the dispersion by means of the droplet size distribution function $u = u(t, y)$ at time t (per unit mass), y being the mass (or volume) of a droplet. By $y_0 \in (0, \infty)$ we denote the maximal droplet mass, which we assume to be a priori known, so that $(0, y_0]$ represents the admissible range of droplet masses. Neglecting dependence on spatial coordinates (for a treatment of

the spatially inhomogeneous case we refer to [28]), the evolution of the system of droplets that undergo both coalescence and breakage can be described by the set of integro-differential equations

$$\begin{aligned} \dot{u}(y) &= \varphi(u)L[u](y), \quad t > 0, \quad y \in (0, y_0], \\ u(0, y) &= u^0(y), \quad y \in (0, y_0], \end{aligned} \quad (*)$$

where u^0 is a given initial distribution. The reaction terms are defined as

$$L[u] := L_b[u] + L_c[u] + L_s[u]$$

whereby for $y \in (0, y_0]$

$$\begin{aligned} L_b[u](y) &:= \int_y^{y_0} \gamma(y', y)u(y') \, dy' - \frac{1}{2}u(y) \int_0^y \gamma(y, y') \, dy', \\ L_c[u](y) &:= \frac{1}{2} \int_0^y K(y', y - y')P(y', y - y')u(y')u(y - y') \, dy' \\ &\quad + \frac{1}{2} \int_y^{y_0} \int_0^{y'} K(y'', y' - y'')Q(y'', y' - y'') \\ &\quad \quad \quad \beta_c(y', y)u(y'')u(y' - y'') \, dy''dy' \\ &\quad - u(y) \int_0^{y_0-y} K(y, y')\{P(y, y') + Q(y, y')\}u(y') \, dy', \\ L_s[u](y) &:= \frac{1}{2} \int_{y_0}^{y_0+y} \int_{y'-y_0}^{y_0} K(y'', y' - y'')\beta_s(y', y)u(y'')u(y' - y'') \, dy''dy' \\ &\quad - u(y) \int_{y_0-y}^{y_0} K(y, y')u(y') \, dy'. \end{aligned}$$

The linear operator $L_b[u]$ gives the gain and loss of droplets of mass y due to binary breakage, where the kernel $\gamma(y, y')$ represents the rate at which a droplet of mass y decays into a droplet of mass $y' \in (0, y)$. Binary breakage in particular means

$$\gamma(y, y') = \gamma(y, y - y'), \quad 0 < y' < y \leq y_0. \quad (1.1)$$

When two droplets y and y' with cumulative mass $y + y' \leq y_0$ collide, three different events may arise being described by the collision operator $L_c[u]$. They either coalesce with probability $P(y, y')$, or a shattering of these droplets occurs with probability $Q(y, y')$, or just nothing happens meaning that the droplets remain unchanged. Obviously, it then holds

$$0 \leq P(y, y') + Q(y, y') \leq 1, \quad 0 < y + y' \leq y_0. \quad (1.2)$$

The symmetric function $K(y, y')$ denotes the rate of binary collision. Further, $\beta_c(y + y', y'')$ is the distribution function of products from a particle $y + y' \in (0, y_0]$ shattering after collision, and β_c satisfies

$$\beta_c(y + y', y'') = \beta_c(y + y', y + y' - y''), \quad 0 < y'' < y + y' \leq y_0. \quad (1.3)$$

The factors $1/2$ come in to compensate for double counting.

The scattering operator $L_s[u]$ represents the interaction of two colliding droplets whose cumulative mass exceeds y_0 and immediately split into two droplets both with mass in $(0, y_0]$. The distribution function $\beta_s(y + y', y'')$ for $y + y' \in (y_0, 2y_0]$ has an analogue meaning as $\beta_c(y + y', y'')$ for $y + y' \in (0, y_0]$ above. Therefore

$$\beta_s(y + y', y'') = \beta_s(y + y', y + y' - y'') , \quad 0 < y + y' - y_0 \leq y'' \leq y_0 . \quad (1.4)$$

We assume that β_c and β_s merely depend on the cumulative mass $y + y'$ of the colliding droplets although there would barely be a difference in the further analysis to allow a dependence on each colliding droplet.

Finally, the efficiency factor $\varphi(u)$ linked to some average properties of the dispersion enhances or depresses the dynamics while the mechanical structure of the interactions is described by the kernels $\gamma, \beta_c, \beta_s, K, P$, and Q . For instance, $\varphi(u)$ may be of the form

$$\varphi(u) = \Phi \left(\int_0^{y_0} u(y) \, dy , \int_0^{y_0} y^{2/3} u(y) \, dy \right) , \quad (1.5)$$

where $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}^+$ is a given function. This means that $\varphi(u)$ is related to the total number of droplets and the total surface area. Clearly, no mathematical substantial differences arise if one considers for each process an individual efficiency factor. But to keep the notation simple, we omit this.

The model considered in [4], [13], [14], [21] can be recovered from (*) by putting $P \equiv 1$. In particular, the shattering terms then drop since $Q \equiv 0$ according to (1.2). For these reduced equations global existence and uniqueness of non-negative and mass preserving solutions is shown in [13], which are Lipschitz continuous with respect to droplet size. These results are extended in [4] to include breakage kernels with singularities. Numerical simulations are performed in [21] exhibiting some interesting features concerning the qualitative behaviour of the solutions for large times.

Finally, a slightly modified version of model (*) — including also multiple breakage — is considered by the author [27]. In the particular case of binary breakage, solutions belonging to the space $L_1((0, y_0])$ are shown to exist globally in time and to be unique.

It is the purpose of the present paper to investigate the long-time behaviour of the particular solutions of [27] assuming that the processes under consideration are somehow reversible. More precisely, we assume that the kernels satisfy an extended version of the so-called *detailed balance condition* (see hypothesis (H_6) below) guaranteeing the existence of equilibria and also providing a Lyapunov function. Such a reversibility condition on the kernels was used in various papers in order to study the qualitative behaviour of solutions for large times. For a treatment of the asymptotic behaviour of solutions to the discrete analogue of (*), we refer to [5], [6], [10] concerning the spatially homogeneous case and to [9], [18] for the case including diffusion (see also [2], [19] for the Becker-Döring equations). Asymptotics for the continuous model without diffusion is studied in [1], [17], [23], [24] whereas the long-time behaviour for continuous coagulation-fragmentation models taking into account diffusion is investigated in [16]. Note that all of the just cited papers consider neither the possibility of shattering nor the existence of a maximal droplet

mass so that there is also no scattering. In this article we include both of these processes. Inspired by the work of [16], we prove in section 2 that the solutions converge (with respect to the L_1 -topology) towards the unique equilibrium with the same mass as the initial distribution. Moreover, in section 3 we derive stability of these equilibria in a suitable topology.

2. Trend to Equilibrium

In the sequel, we put $L_1 := L_1((0, y_0])$ and denote by $|\cdot|_1$ the norm of L_1 . The closed subset of L_1 consisting of all $u \in L_1$ which are non-negative almost everywhere is denoted by L_1^+ . Furthermore, $L_{1,w}$ stands for the space L_1 endowed with its weak topology.

Throughout this article we assume that the following hypotheses are satisfied:

(H₁) $\varphi : L_1 \rightarrow (0, \infty)$ is uniformly Lipschitz continuous on bounded sets, weakly sequentially continuous, and bounded;

(H₂) γ is a measurable function from $\Delta := \{(y, y') ; 0 < y' < y \leq y_0\}$ into \mathbb{R}^+ satisfying (1.1), and there exists $m_\gamma > 0$ with

$$\int_0^y \gamma(y, y') \, dy' \leq m_\gamma, \quad \text{a.a. } y \in (0, y_0] ;$$

(H₃) β_c is a measurable function from Δ into \mathbb{R}^+ satisfying (1.3) and

$$\int_0^{y+y'} \beta_c(y+y', y'') \, dy'' = 2, \quad (2.1)$$

for a.a. $(y, y') \in (0, y_0]^2$ with $y+y' \in (0, y_0]$;

(H₄) β_s is a measurable function from $\{(y, y') ; 0 < y - y_0 \leq y' \leq y_0\}$ into \mathbb{R}^+ satisfying (1.4) and

$$\int_{y+y'-y_0}^{y_0} \beta_s(y+y', y'') \, dy'' = 2, \quad (2.2)$$

for a.a. $(y, y') \in (0, y_0]^2$ with $y+y' \in (y_0, 2y_0]$;

(H₅) $P, Q, K \in L_\infty((0, y_0]^2, \mathbb{R}^+)$ are symmetric and P, Q satisfy (1.2) whereas $PK > 0$ a.e.;

(H₆) there exists $H \in L_1^+$ with $\text{ess-inf } H > 0$ and

(i) for $0 < y + y' < y_0$ it holds

$$\gamma(y+y', y)H(y+y') = P(y, y')K(y, y')H(y)H(y'),$$

(ii) for $0 < y + y', y + y'' < y_0$ it holds

$$\begin{aligned} \beta_c(y, y')Q(y'', y-y'')K(y'', y-y'')H(y'')H(y-y'') \\ = \beta_c(y, y'')Q(y', y-y')K(y', y-y')H(y')H(y-y'), \end{aligned}$$

(iii) for $0 < y - y_0 < y'$, $y'' < y_0$ it holds

$$\begin{aligned} \beta_s(y, y')K(y'', y - y'')H(y'')H(y - y'') \\ = \beta_s(y, y'')K(y', y - y')H(y')H(y - y') . \end{aligned}$$

We refer to Examples 2.12 for kernels satisfying the hypotheses above. Equalities (2.1) and (2.2) reflect binary breakage in the shattering and scattering processes, respectively. Observe that in combination with (1.3) and (1.4) they additionally imply

$$\int_0^{y+y'} y'' \beta_c(y + y', y'') \, dy'' = y + y' , \quad (2.3)$$

for a.a. $(y, y') \in (0, y_0]^2$ with $y + y' \in (0, y_0]$ and

$$\int_{y+y'-y_0}^{y_0} y'' \beta_s(y + y', y'') \, dy'' = y + y' , \quad (2.4)$$

for a.a. $(y, y') \in (0, y_0]^2$ with $y + y' \in (y_0, 2y_0]$. In other words, shattering and scattering are mass preserving processes.

Before making use of hypothesis (H_6) , let us collect some already proven facts on global existence of solutions to problem $(*)$, that is, for the ordinary differential equation

$$\begin{aligned} \dot{u} &= \varphi(u)L[u] , \quad t > 0 , \\ u(0) &= u^0 , \end{aligned} \quad (**)$$

considered in L_1 .

Theorem 2.1. *Suppose that hypotheses $(H_1) - (H_5)$ are satisfied. Then, given any $u^0 \in L_1^+$, problem $(**)$ admits a unique solution $u(\cdot; u^0) \in C^1(\mathbb{R}^+, L_1)$ which, in addition, is non-negative and preserves the total mass, i.e.*

$$\int_0^{y_0} y u(t; u^0)(y) \, dy = \int_0^{y_0} y u^0(y) \, dy , \quad t \geq 0 .$$

Moreover, the map $(t, u^0) \mapsto u(t; u^0)$ defines a semiflow on L_1^+ .

Proof. This follows by an obvious modification of the proofs in [27] (there, the case $Q \equiv 1 - P$ is treated). A detailed proof is also given in [28]. \square

In the following, we denote by $u := u(\cdot; u^0) \in C^1(\mathbb{R}^+, L_1)$ the unique solution to $(**)$, and we write

$$u(t, y) := u(t; u^0)(y) , \quad \text{a.a. } y \in (0, y_0] , \quad t \geq 0 ,$$

if the initial value $u^0 \in L_1^+$ is fixed. Sometimes we suppress any of the variables t and y in a given formula. Further, c or $c(u^0)$ will denote various constants, which may differ from occurrence to occurrence, but which are always independent of the free variables.

It is an easy consequence of hypothesis (H_6) that the function $u_\alpha \in L_1^+$, given by

$$u_\alpha(y) := H(y)e^{\alpha y} , \quad \text{a.a. } y \in (0, y_0] , \quad (2.5)$$

is for each $\alpha \in \mathbb{R}$ an equilibrium of problem (**). Let us then introduce the map $V : L_1^+ \rightarrow \mathbb{R}^+ \cup \{\infty\}$ according to

$$V(v) := \int_0^{y_0} \left\{ v(y) \left[\log \frac{v(y)}{H(y)} - 1 \right] + H(y) \right\} dy, \quad v \in L_1^+,$$

which will turn out to be a Lyapunov function for (**). Note that Fatou's lemma entails that V is sequentially lower semi-continuous. Hence V is weakly sequentially lower semi-continuous due to its convexity (see [12, Prop.2.3]).

Next, the proof of [16, Lem.3.1] can easily be modified to yield the following lemma, which will guarantee that the orbit of the motion through $u^0 \in L_1^+$ is relatively weakly compact in L_1 provided $V(u^0) < \infty$.

Lemma 2.2. *Let $w \in L_1^+$ be such that $V(w) < \infty$. Then, for each $\alpha \geq e^2$ and each measurable subset A of $(0, y_0]$, it holds*

$$\int_A w(y) dy \leq 2\alpha \int_A H(y) dy + \frac{2}{\log \alpha} V(w).$$

Further, it is not difficult to adapt the ideas of [16, Lem.C.1] in order to prove the next result. We refrain from giving details and refer to [28, Lem.3.9].

Lemma 2.3. *Suppose that $w \in L_1^+$ satisfies*

$$\gamma(y + y', y)w(y + y') = P(y, y')K(y, y')w(y)w(y') \quad (2.6)$$

for a.a. $(y, y') \in (0, y_0]^2$ with $0 < y + y' \leq y_0$. Then either $w = 0$ a.e. or there exists $\alpha \in \mathbb{R}$ such that $w(y) = H(y)e^{\alpha y}$ for a.a. $y \in (0, y_0]$.

The main ingredient for examining large-time behaviour of the solutions consists of proving that V is a Lyapunov function for (**), that is, that V is decreasing along orbits. Such a result will make heavily use of hypothesis (H_6) as well as of formulas (2.1) and (2.2). In order to carry through rigorously the technical details, we need an upper and a lower bound for the solutions. This may be obtained by approximating the solution to (**) by solutions to a modified problem, where the initial value and the kernels are truncated in a suitable way paying attention to the detailed balance condition (H_6) . But then these truncated kernels do no longer obey equalities of type (2.1) and (2.2). Hence, we also have to alter the reaction terms slightly in order to guarantee that V is still decreasing along orbits of solutions to the modified problem. For that purpose, let us introduce some further notations. Define the set

$$\mathcal{E} := \{(y, y') \in (0, y_0]^2; y + y' < y_0\},$$

as well as for $n \geq 1$ the sets

$$\begin{aligned} A_n &:= \{(y, y') \in \mathcal{E}; \gamma(y + y', y) \leq n\}, \\ B_n &:= \{(y, y') \in \mathcal{E}; \beta_c(y + y', y) \leq n\}, \\ C_n &:= \{(y, y') \in (0, y_0]^2 \setminus \mathcal{E}; \beta_s(y + y', y) \leq n\}, \end{aligned}$$

and observe that (y, y') belongs to any one of the sets A_n, B_n , or C_n if and only if (y', y) does. Further, truncate the kernels according to

$$\begin{aligned}\gamma_n(y + y', y) &:= \begin{cases} \gamma(y + y', y), & (y, y') \in A_n \cap B_n, \\ 0, & \text{else,} \end{cases} \\ \beta_{c,n}(y + y', y) &:= \begin{cases} \beta_c(y + y', y), & (y, y') \in A_n \cap B_n, \\ 0, & \text{else,} \end{cases} \\ \beta_{s,n}(y + y', y) &:= \begin{cases} \beta_s(y + y', y), & (y, y') \in C_n, \\ 0, & \text{else,} \end{cases} \\ K_n(y, y') &:= \begin{cases} K(y, y'), & (y, y') \in (A_n \cap B_n) \cup C_n, \\ 0, & \text{else.} \end{cases}\end{aligned}$$

Then K_n is symmetric and $\gamma_n, \beta_{c,n}$, and $\beta_{s,n}$ satisfy hypotheses $(H_2), (H_3)$, and (H_4) , respectively. Furthermore,

$$\gamma_n \nearrow \gamma, \quad \beta_{c,n} \nearrow \beta_c, \quad \beta_{s,n} \nearrow \beta_s, \quad K_n \nearrow K, \quad (2.7)$$

pointwise on the domains of γ, β_c, β_s , and K . Finally, the truncated kernels satisfy the detailed balance condition (H_6) with the same function H and the same probabilities P and Q .

In addition, define for $w \in L_1$ and a.a. $y \in (0, y_0]$

$$\begin{aligned}L_{b,n}[w](y) &:= \int_y^{y_0} \gamma_n(y', y) w(y') \, dy' - \frac{1}{2} w(y) \int_0^y \gamma_n(y, y') \, dy', \\ L_{c,n}[w](y) &:= \frac{1}{2} \int_0^y K_n(y', y - y') P(y', y - y') w(y') w(y - y') \, dy' \\ &\quad + \frac{1}{2} \int_y^{y_0} \int_0^{y'} K_n(y'', y' - y'') Q(y'', y' - y'') \\ &\quad \quad \quad \beta_{c,n}(y', y) w(y'') w(y' - y'') \, dy'' dy' \\ &\quad - w(y) \int_0^{y_0 - y} K_n(y, y') P(y, y') w(y') \, dy' \\ &\quad - \frac{1}{2} w(y) \int_0^{y_0 - y} \int_0^{y + y'} \beta_{c,n}(y + y', y'') \, dy'' K_n(y, y') Q(y, y') w(y') \, dy', \\ L_{s,n}[w](y) &:= \frac{1}{2} \int_{y_0}^{y_0 + y} \int_{y' - y_0}^{y_0} \beta_{s,n}(y', y) K_n(y'', y' - y'') w(y'') w(y' - y'') \, dy'' dy' \\ &\quad - \frac{1}{2} w(y) \int_{y_0 - y}^{y_0} \int_{y + y' - y_0}^{y_0} \beta_{s,n}(y + y', y'') \, dy'' K_n(y, y') w(y') \, dy',\end{aligned}$$

and further

$$L_n[w] := L_{b,n}[w] + L_{c,n}[w] + L_{s,n}[w], \quad w \in L_1.$$

In the sequel, we denote by $|\cdot|_\infty$ the norm of $L_\infty := L_\infty((0, y_0])$.

Lemma 2.4. *Given $n \geq 1$ and any non-negative $w^0 \in L_\infty$, there exists a unique solution $w := w(\cdot; w^0) \in C^1(\mathbb{R}^+, L_\infty)$ for the problem*

$$\begin{aligned} \dot{w} &= \varphi(w)L_n[w] , \quad t > 0 , \\ w(0) &= w^0 . \end{aligned}$$

Moreover, this solution is non-negative and, in addition, if $w^0 \geq r_0$ a.e. for some $r_0 \in (0, \infty)$ then, for any $T > 0$, there exists $r_T > 0$ such that

$$w(t) \geq r_T \quad \text{a.e.} , \quad 0 \leq t \leq T . \quad (2.8)$$

Proof. According to hypotheses $(H_1) - (H_5)$ it holds

$$|\varphi(w)L_n[w]|_\infty \leq c(1 + |w|_1)|w|_\infty , \quad w \in L_\infty . \quad (2.9)$$

From this, existence of a unique solution $w \in C^1(J(w^0), L_\infty)$ follows, where $J(w^0)$ denotes the maximal interval of existence. That this solution is non-negative may be obtained along the lines of the proof of [27, Thm.2.4]. Observe then that

$$\int_0^{y_0} L_{b,n}[v](y) dy \leq c|v|_1 , \quad \int_0^{y_0} L_{c,n}[v](y) dy \leq 0 , \quad \int_0^{y_0} L_{s,n}[v](y) dy = 0 ,$$

for $v \in L_1^+$. Since $w(t) \in L_1^+$ for $t \in J(w^0)$, Gronwall's inequality applies to provide $c := c(w^0)$ with

$$|w(t)|_1 \leq ce^{ct} , \quad t \in J(w^0) ,$$

so that (2.9) entails $J(w^0) = \mathbb{R}^+$. Finally, it remains to prove (2.8). Fix $T > 0$ arbitrarily and put

$$\omega := \|\varphi\|_\infty (m_\gamma + \|K\|_\infty \max_{0 \leq t \leq T} |w(t)|_1) .$$

Since $w(s) \geq 0$ a.e. we deduce for $0 \leq t \leq T$

$$\begin{aligned} w(t) &= e^{-\omega t} w^0 + \int_0^t e^{-\omega(t-s)} \left\{ \varphi(w(s))L_n[w(s)] + \omega w(s) \right\} ds \\ &\geq e^{-\omega T} r_0 =: r_T \quad \text{a.e.} . \end{aligned}$$

□

We also need the following lemma whose prove can be found in [16, Lem.A.2].

Lemma 2.5. *Let $\Omega \subset \mathbb{R}^m$, $m \geq 1$, be a measurable and bounded set. Assume that $h_n, h \in L_\infty(\Omega)$ are such that $\|h_n\|_\infty \leq c$ for $n \geq 1$ and $h_n \rightarrow h$ a.e.. Then, provided $v_n \rightarrow v$ in $L_{1,w}(\Omega)$, it holds $h_n v_n \rightarrow h v$ in $L_{1,w}(\Omega)$.*

The next lemma will ensure in particular that the solutions to the modified problem, being provided by Lemma 2.4, indeed approximate the original solution $u(\cdot; u^0)$.

Lemma 2.6. Assume that $w_n \rightarrow w$ in $L_{1,w}$.

(i) Defining for $(y, y') \in \mathcal{E}$

$$v_n(y, y') := \gamma_n(y + y', y)w_n(y + y') ,$$

and

$$v(y, y') := \gamma(y + y', y)w(y + y') ,$$

it holds $v_n \rightarrow v$ in $L_{1,w}(\mathcal{E})$.

(ii) Defining for $(y, y') \in \mathcal{E}$

$$z_n(y, y') := P(y, y')K_n(y, y')w_n(y)w_n(y')$$

and

$$z(y, y') := P(y, y')K(y, y')w(y)w(y') ,$$

it holds $z_n \rightarrow z$ in $L_{1,w}(\mathcal{E})$.

(iii) It holds $L_n[w_n] \rightarrow L[w]$ in $L_{1,w}$.

Proof. Given $f \in L_\infty(\mathcal{E})$ use Fubini's theorem to deduce

$$\begin{aligned} & \left| \int_{\mathcal{E}} f(y, y') [v_n(y, y') - v(y, y')] \, d(y, y') \right| \\ & \leq \|f\|_\infty \int_0^{y_0} a_n(y) |w(y)| \, dy + \left| \int_0^{y_0} h_n(y) [w(y) - w_n(y)] \, dy \right| \end{aligned} \quad (2.10)$$

where

$$a_n(y) := \int_0^y |\gamma_n(y, y') - \gamma(y, y')| \, dy' , \quad h_n(y) := \int_0^y f(y', y - y') \gamma_n(y, y') \, dy' .$$

Due to hypothesis (H_2) and (2.7), an application of Lebesgue's theorem yields that the first term on the right hand side of (2.10) converges to 0 as $n \rightarrow \infty$. Next observe that for a.a. $y \in (0, y_0]$ we have, in virtue of Fubini's theorem,

$$f(\cdot, y - \cdot) \in L_\infty((0, y)) \quad \text{with} \quad \|f(\cdot, y - \cdot)\|_{L_\infty((0, y))} \leq \|f\|_\infty .$$

We obtain $|h_n|_\infty \leq \|f\|_\infty m_\gamma$ and, using Lebesgue's theorem,

$$h_n(y) \rightarrow h(y) := \int_0^y f(y', y - y') \gamma(y, y') \, dy' , \quad \text{a.a. } y \in (0, y_0] ,$$

where $h \in L_\infty$. Lemma 2.5 entails now $v_n \rightarrow v$ in $L_{1,w}(\mathcal{E})$. All other statements can be proven in a similar way (for (iii) recall (2.1) and (2.2)). Therefore, we refrain from giving more details and refer to [28]. \square

Let us introduce some further notations. Define the map $\mathcal{J} : \mathbb{R}^2 \rightarrow \mathbb{R}^+ \cup \{\infty\}$ by

$$\mathcal{J}(a, b) := \begin{cases} (a - b)(\log a - \log b) , & a, b > 0 , \\ 0 , & a = b = 0 , \\ \infty , & \text{else} . \end{cases}$$

In order to shorten the formulas we agree upon putting

$$y''' \equiv y + y' - y'' , \quad 0 < y'' < y + y' .$$

Moreover, we set for $v \in L_1^+$

$$\begin{aligned} D(v) &:= \frac{1}{2} \int_{\mathcal{E}} \mathcal{J} \left(P(y, y') K(y, y') v(y) v(y') , \gamma(y + y', y) v(y + y') \right) d(y, y') , \\ F(v) &:= \frac{1}{8} \int_{\mathcal{W}} \mathcal{J} \left(\beta_c(y + y', y) Q(y'', y''') K(y'', y''') v(y'') v(y''') , \right. \\ &\quad \left. \beta_c(y + y', y'') Q(y, y') K(y, y') v(y) v(y') \right) d(y, y', y'') , \\ G(v) &:= \frac{1}{8} \int_{\mathcal{S}} \mathcal{J} \left(\beta_s(y + y', y) K(y'', y''') v(y'') v(y''') , \right. \\ &\quad \left. \beta_s(y + y', y'') K(y, y') v(y) v(y') \right) d(y, y', y'') , \end{aligned}$$

where the sets \mathcal{W} and \mathcal{S} are given by

$$\begin{aligned} \mathcal{W} &:= \{ (y, y', y'') \in (0, y_0]^3 ; y'' < y + y' < y_0 \} , \\ \mathcal{S} &:= \{ (y, y', y'') \in (0, y_0]^3 ; y_0 - y'' < y + y' - y'' < y_0 \} . \end{aligned}$$

Finally, we define $D_n(v)$, $F_n(v)$, and $G_n(v)$ analogously but with $(\gamma_n, \beta_{c,n}, \beta_{s,n}, K_n)$ instead of $(\gamma, \beta_c, \beta_s, K)$.

Now we are in position to prove that V is indeed a Lyapunov function for (**).

Proposition 2.7. *Let $u^0 \in L_1^+$ be such that $V(u^0) < \infty$ and denote by $u = u(\cdot; u^0)$ the unique, non-negative solution to (**) in $C^1(\mathbb{R}^+, L_1)$. Then it holds*

$$0 \leq V(u(t)) \leq V(u(s)) < \infty , \quad t \geq s \geq 0 , \quad (2.11)$$

and

$$[t \mapsto \varphi(u(t)) D(u(t))] \in L_1(\mathbb{R}^+) . \quad (2.12)$$

Proof. For $n \geq 1$ set

$$u_n^0(y) := \min \{ n, \max \{ u^0(y), H(y)/n \} \} , \quad \text{a.a. } y \in (0, y_0) ,$$

and observe that $0 < \min \{ n, \frac{1}{n} \text{ess-inf } H \} \leq u_n^0 \leq n$ a.e. and $u_n^0 \rightarrow u^0$ in L_1 . Further we have

$$\int_0^{y_0} u_n^0 \log \frac{u_n^0}{H} dy \leq \left(\int_{S_n} + \int_{T_n} \right) u^0 \log \frac{u^0}{H} dy , \quad n \geq 1 ,$$

where we put

$$S_n := \left[\frac{H}{n} \leq u^0 < n \right] \quad \text{and} \quad T_n := [H < n \leq u^0] .$$

Taking into account that $V(u^0) < \infty$ and $r |\log r| \leq r \log r + \frac{2}{\varepsilon}$, $r \geq 0$, imply $u^0 \log \frac{u^0}{H} \in L_1$, Lebesgue's theorem yields

$$\limsup_n V(u_n^0) \leq V(u^0) . \quad (2.13)$$

Next, Lemma 2.4 entails the existence of a solution $u_n := u_n(\cdot; u_n^0) \in C^1(\mathbb{R}^+, L_\infty)$ to the problem

$$\begin{aligned} \dot{w} &= \varphi(w)L_n[w], \quad t > 0, \\ w(0) &= u_n^0 \end{aligned}$$

satisfying for each $T > 0$

$$0 < r_n^1(T) \leq u_n(t) \leq r_n^2(T) < \infty \quad \text{a.e.}, \quad 0 \leq t \leq T, \quad (2.14)$$

for some constants $r_n^j(T)$. This enables us to deduce

$$\frac{d}{dt}V(u_n(t)) = \varphi(u_n(t)) \int_0^{y_0} \log \frac{u_n(t, y)}{H(y)} L_n[u_n(t)](y) dy \quad (2.15)$$

for $n \geq 1$ and $0 \leq t \leq T$. Note that Fubini's theorem applies throughout in the following because of (2.14). Little effort then yields

$$\int_0^{y_0} \log \frac{u_n(t, y)}{H(y)} \left\{ L_{b,n}[u_n(t)](y) + L_{c,n}^{(P)}[u_n(t)](y) \right\} dy = -D_n(u_n(t)), \quad (2.16)$$

for $n \geq 1$ and $0 \leq t \leq T$, where $L_{c,n}^{(P)}$ consists of those integral terms of $L_{c,n}$ involving P but not Q . Further we compute

$$\begin{aligned} & \int_0^{y_0} \log \frac{u_n(y)}{H(y)} L_{s,n}[u_n](y) dy \\ &= \frac{1}{2} \int_{\mathcal{S}} \left\{ \log \frac{u_n(y'')}{H(y'')} - \log \frac{u_n(y)}{H(y)} \right\} \\ & \quad \beta_{s,n}(y + y', y'') K_n(y, y') u_n(y) u_n(y') d(y, y', y'') \\ &= \frac{1}{4} \int_{\mathcal{S}} \left\{ \log \frac{u_n(y'') u_n(y''')}{H(y'') H(y''')} - \log \frac{u_n(y) u_n(y')}{H(y) H(y')} \right\} \\ & \quad \beta_{s,n}(y + y', y'') K_n(y, y') u_n(y) u_n(y') d(y, y', y''), \end{aligned}$$

where we have taken into account the symmetry of K_n and that $\beta_{s,n}$ satisfies (1.4). The transformation $\mathcal{S} \rightarrow \mathcal{S}$, $(y, y', y'') \mapsto (y'', y''', y)$ entails then that the right hand side of the above equality coincides with

$$\begin{aligned} & \frac{1}{8} \int_{\mathcal{S}} \left\{ \log \frac{u_n(y'') u_n(y''')}{H(y'') H(y''')} - \log \frac{u_n(y) u_n(y')}{H(y) H(y')} \right\} \\ & \quad \beta_{s,n}(y + y', y'') K_n(y, y') u_n(y) u_n(y') d(y, y', y'') \\ & + \frac{1}{8} \int_{\mathcal{S}} \left\{ \log \frac{u_n(y) u_n(y')}{H(y) H(y')} - \log \frac{u_n(y'') u_n(y''')}{H(y'') H(y''')} \right\} \\ & \quad \beta_{s,n}(y + y', y) K_n(y'', y''') u_n(y'') u_n(y''') d(y, y', y''). \end{aligned}$$

Finally, due to hypothesis (H_6) we may rewrite this last expression to get

$$\int_0^{y_0} \log \frac{u_n(t, y)}{H(y)} L_{s,n}[u_n(t)](y) dy = -G_n(u_n(t)), \quad (2.17)$$

for $n \geq 1$ and $0 \leq t \leq T$. Likewise one derives

$$\int_0^{y_0} \log \frac{u_n(t, y)}{H(y)} L_{c,n}^{(Q)}[u_n(t)](y) \, dy = -F_n(u_n(t)) , \quad (2.18)$$

where $L_{c,n}^{(Q)}$ are those integral terms of $L_{c,n}$ involving Q but not P . Therefore, (2.15)-(2.18) in combination with (2.13) yield for $n \geq 1$ and $0 \leq t \leq T$

$$\begin{aligned} V(u_n(t)) + \int_0^t \varphi(u_n(\sigma)) \{D_n(u_n(\sigma)) + F_n(u_n(\sigma)) + G_n(u_n(\sigma))\} \, d\sigma \\ = V(u_n^0) \leq c(u^0) < \infty . \end{aligned} \quad (2.19)$$

Consequently,

$$V(u_n(t)) \leq c(u^0) , \quad n \geq 1 , \quad t \geq 0 , \quad (2.20)$$

since each of the terms $D_n(u_n(\sigma))$, $F_n(u_n(\sigma))$, and $G_n(u_n(\sigma))$ is non-negative. Hence, Lemma 2.2 leads to

$$|u_n(t)|_1 \leq c(u^0) , \quad n \geq 1 , \quad t \geq 0 , \quad (2.21)$$

and invoking additionally the Dunford-Pettis theorem [11, Thm.4.21.2] we see that the set $\{u_n(t); n \geq 1\}$ is relatively weakly compact in L_1 for each $t \geq 0$. Next, from (2.7) and hypotheses $(H_1) - (H_5)$ we derive

$$|\varphi(v)L_n[v]|_1 \leq c(1 + |v|_1)|v|_1 , \quad v \in L_1 , \quad n \geq 1 , \quad (2.22)$$

with c being independent of $n \geq 1$. This and (2.21) imply

$$|u_n(t) - u_n(s)|_1 \leq c(u^0)|t - s| , \quad t, s \geq 0 , \quad n \geq 1 . \quad (2.23)$$

In particular, the set $\{u_n; n \geq 1\}$ is equicontinuous with respect to the weak topology of L_1 . Now fix $T > 0$ arbitrarily. Then the Arzelà-Ascoli theorem [26, Thm.1.3.2] entails that there exist $\bar{u} \in C([0, T], L_{1,w})$ and a subsequence (n') such that

$$u_{n'} \rightarrow \bar{u} \quad \text{in } C([0, T], L_{1,w}) . \quad (2.24)$$

Clearly, \bar{u} belongs to $C^{1-}([0, T], L_1)$ due to (2.23), that is, \bar{u} is Lipschitz continuous with respect to the L_1 -topology. Further, thanks to Lemma 2.6 we have

$$L_{n'}[u_{n'}(\sigma)] \rightarrow L[\bar{u}(\sigma)] \quad \text{in } L_{1,w} , \quad 0 \leq \sigma \leq T . \quad (2.25)$$

Since φ is weakly sequentially continuous, an application of Lebesgue's theorem, (2.21), (2.22), (2.24), and (2.25) yield

$$\int_0^t \varphi(u_{n'}(\sigma)) L_{n'}[u_{n'}(\sigma)] \, d\sigma \longrightarrow \int_0^t \varphi(\bar{u}(\sigma)) L[\bar{u}(\sigma)] \, d\sigma \quad \text{in } L_{1,w} , \quad 0 \leq t \leq T ,$$

so that a renewed use of (2.24) shows

$$\bar{u}(t) = u^0 + \int_0^t \varphi(\bar{u}(\sigma)) L[\bar{u}(\sigma)] \, d\sigma , \quad 0 \leq t \leq T .$$

Hence $\bar{u} = u(\cdot; u^0)|_{[0, T]}$ due to uniqueness of solutions to (**). Consequently we have

$$u_{n'} \rightarrow u(\cdot; u^0) \quad \text{in } C([0, T], L_{1, w}) . \quad (2.26)$$

Since V is weakly lower semi-continuous and since $T > 0$ was arbitrary, we deduce from (2.19) and (2.13) that (2.11) is indeed true for $t \geq s = 0$. The semiflow property then yields the general case of (2.11).

Hence it remains to prove (2.12). According to (2.26) we may apply Lemma 2.6 to obtain

$$\begin{aligned} \gamma_{n'}(y + y', y) u_{n'}(\sigma, y + y') &\rightarrow \gamma(y + y', y) u(\sigma, y + y') \quad \text{in } L_{1, w}(\mathcal{E}) , \\ P(y, y') K_{n'}(y, y') u_{n'}(\sigma, y) u_{n'}(\sigma, y') &\rightarrow P(y, y') K(y, y') u(\sigma, y) u(\sigma, y') \quad \text{in } L_{1, w}(\mathcal{E}) , \end{aligned}$$

for $0 \leq \sigma \leq T$. Since the function \mathcal{J} , appearing in the definition of $D(v)$, is convex and lower semi-continuous, we obtain from the above convergence, from Fatou's lemma, and from (2.19) that

$$\int_0^T \varphi(u(\sigma)) D(u(\sigma)) \, d\sigma \leq \liminf_{n'} \int_0^T \varphi(u_{n'}(\sigma)) D_{n'}(u_{n'}(\sigma)) \, d\sigma \leq c(u^0) ,$$

whereby $c(u^0)$ does not depend on $T > 0$. □

Recall that the equilibria u_α , $\alpha \in \mathbb{R}$, are given by (2.5). Clearly, given any $\varrho > 0$ there exists $\alpha(\varrho) \in \mathbb{R}$ uniquely such that $M(u_{\alpha(\varrho)}) = \varrho$, where the mass $M(v)$ of $v \in L_1^+$ is defined as

$$M(v) := \int_0^{y_0} y v(y) \, dy .$$

Now we can state the result concerning convergence towards equilibrium.

Theorem 2.8. *Given $u^0 \in L_1^+ \setminus \{0\}$ with $V(u^0) < \infty$ choose $\alpha \in \mathbb{R}$ such that $M(u_\alpha) = M(u^0)$. Then, given any sequence $t_n \nearrow \infty$ and any $T > 0$, the solution $u = u(\cdot; u^0)$ to problem (**) satisfies*

$$u(\cdot + t_n; u^0) \rightarrow u_\alpha \quad \text{in } C([0, T], L_{1, w}) . \quad (2.27)$$

In addition, if there exists $r \in L_1^+$ such that for a.a. $y \in (0, y_0)$

$$\gamma(\cdot, y) \leq r(y) \quad \text{a.e. on } (y, y_0) \quad (2.28)$$

and if $u^0 > 0$ a.e., then

$$u(\cdot + t_n; u^0) \rightarrow u_\alpha \quad \text{in } C([0, T], L_1) . \quad (2.29)$$

Proof. Put

$$u_n(t) := u(t + t_n; u^0) = u(t; u(t_n; u^0)) , \quad t \geq 0 , \quad n \geq 1 ,$$

so that, according to Proposition 2.7,

$$V(u_n(t)) \leq V(u^0) , \quad t \geq 0 , \quad n \geq 1 . \quad (2.30)$$

Analogously to the proof of Proposition 2.7 we deduce the existence of a function $\bar{u} \in C^{1-}([0, T], L_1)$ and of a subsequence (n') such that $u_{n'} \rightarrow \bar{u}$ in $C([0, T], L_{1,w})$. Obviously, we have $\bar{u}(t) \in L_1^+$ for $0 \leq t \leq T$. Further, as in the proof of Proposition 2.7 we infer

$$0 \leq \int_0^T \varphi(\bar{u}(t))D(\bar{u}(t)) dt \leq \liminf_{n'} \int_0^T \varphi(u_{n'}(t))D(u_{n'}(t)) dt .$$

Thanks to (2.12) the latter expression equals zero. Therefore, $D(\bar{u}(t)) = 0$ for a.a. $0 \leq t \leq T$ since φ has no zeros. By definition of D , Lemma 2.3 entails that $\bar{u}(t)$ is an equilibrium of the form (2.5) for a.a. $t \in [0, T]$. But since

$$M(\bar{u}(t)) = M(u_{n'}(t)) = M(u^0) = M(u_\alpha) , \quad 0 \leq t \leq T ,$$

according to Theorem 2.1, we deduce that \bar{u} is independent of time due to continuity, and it coincides with u_α . Therefore, $u_{n'} \rightarrow u_\alpha$ in $C([0, T], L_{1,w})$, which leads to (2.27) since the limit does not depend on the extracted subsequence.

Let (2.28) be true so that (2.27) implies for $T > 0$

$$L_b^1[u_n(t)](y) \rightarrow L_b^1[u_\alpha](y) , \quad \text{a.a. } y \in (0, y_0] , \quad 0 \leq t \leq T ,$$

where we put

$$L_b^1[v](y) := \int_y^{y_0} \gamma(y', y)v(y') dy' , \quad \text{a.a. } y \in (0, y_0] , \quad v \in L_1 .$$

Moreover, invoking (2.30) and Lemma 2.2 we get

$$|L_b^1[u_n(t)](y)| \leq |u_n(t)|_1 r(y) \leq c(u^0) r(y) , \quad \text{a.a. } y \in (0, y_0] , \quad 0 \leq t \leq T , \quad (2.31)$$

with $c(u^0) > 0$ depending neither on $n \geq 1$ nor on $t \in [0, T]$. Thus, Lebesgue's theorem and (2.27) entail

$$\varphi(u_n)L_b^1[u_n] \rightarrow \varphi(u_\alpha)L_b^1[u_\alpha] \quad \text{in } L_1((0, T) \times (0, y_0]) , \quad (2.32)$$

since φ is weakly sequentially continuous and bounded. For $v \in L_1$ set

$$h(v)(y) := \int_0^{y_0-y} P(y, y')K(y, y')v(y') dy' , \quad \text{a.a. } y \in (0, y_0] .$$

Analogously as above it then holds

$$\varphi(u_n)h(u_n) \rightarrow \varphi(u_\alpha)h(u_\alpha) \quad \text{in } L_1((0, T) \times (0, y_0]) . \quad (2.33)$$

Next, take up the idea of the proof of Lemma 2.4 in order to deduce that $u^0 > 0$ a.e. implies $u(t; u^0) > 0$ a.e. for each $t \geq 0$. Fix $\lambda > 1$ and observe that the inequality

$$|\eta - \xi| \leq (\lambda - 1)\xi + \frac{1}{\log \lambda}(\eta - \xi)(\log \eta - \log \xi) , \quad \xi, \eta > 0 ,$$

holds, from which we derive

$$\begin{aligned} & |\varphi(u_n)u_n h(u_n) - \varphi(u_n)L_b^1[u_n]|_{L_1((0,T) \times (0,y_0))} \\ & \leq \int_0^T \varphi(u_n) \int_0^{y_0} \int_0^{y_0-y} |P(y,y')K(y,y')u_n(y)u_n(y') \\ & \quad - \gamma(y+y',y)u_n(y+y')| \, dy' dy dt \\ & \leq (\lambda - 1) |\varphi(u_n)L_b^1[u_n]|_{L_1((0,T) \times (0,y_0))} + \frac{2}{\log \lambda} \int_0^T \varphi(u_n) D(u_n) \, dt . \end{aligned}$$

Taking the $\limsup_{n \nearrow \infty}$ on both sides and letting then λ tend to 1, (2.12), (2.31), and (2.32) provide

$$\varphi(u_n)u_n h(u_n) \rightarrow \varphi(u_\alpha)L_b^1[u_\alpha] = \varphi(u_\alpha)u_\alpha h(u_\alpha) \quad \text{in } L_1((0,T) \times (0,y_0))$$

as $n \nearrow \infty$, whereby the equality is implied by hypothesis (H_6) . Therefore, recalling (2.33), we may extract a subsequence (n') such that $\varphi(u_{n'})u_{n'}h(u_{n'})$ and $\varphi(u_{n'})h(u_{n'})$ converge pointwise a.e. on $(0,T) \times (0,y_0]$ towards $\varphi(u_\alpha)u_\alpha h(u_\alpha)$ and $\varphi(u_\alpha)h(u_\alpha)$, respectively. But this implies that $u_{n'} \rightarrow u_\alpha$ a.e. on $(0,T) \times (0,y_0]$ since $\varphi(u_\alpha)h(u_\alpha) > 0$ in virtue of hypotheses (H_1) and (H_5) . Analogously to (2.21), the set $\{u_n(t); n \geq 1, 0 \leq t \leq T\}$ is bounded in L_1 so that (2.27) gives $u_{n'} \rightarrow u_\alpha$ in $L_{1,w}((0,T) \times (0,y_0])$ and a.e. on $(0,T) \times (0,y_0]$. Hence

$$u_{n'} \rightarrow u_\alpha \quad \text{in } L_1((0,T) \times (0,y_0)) \quad (2.34)$$

from which we derive (see [27, Lem.2.1])

$$L[u_{n'}] \rightarrow L[u_\alpha] = 0 \quad \text{in } L_1((0,T) \times (0,y_0)) . \quad (2.35)$$

Observing

$$u_{n'}(t) = u_{n'}(s) + \int_s^t \varphi(u_{n'}(\sigma))L[u_{n'}(\sigma)] \, d\sigma , \quad 0 \leq s \leq t ,$$

we then see that for each $t \in (0,T]$

$$\begin{aligned} t|u_{n'}(t) - u_\alpha|_1 & \leq |u_{n'} - u_\alpha|_{L_1((0,T) \times (0,y_0))} + \|\varphi\|_\infty \int_0^t \int_s^t |L[u_{n'}(\sigma)]|_1 \, d\sigma ds \\ & \leq |u_{n'} - u_\alpha|_{L_1((0,T) \times (0,y_0))} + T\|\varphi\|_\infty |L[u_{n'}]|_{L_1((0,T) \times (0,y_0))} . \end{aligned}$$

Recalling (2.34) and (2.35) it therefore holds $u_{n'} \rightarrow u_\alpha$ in $C((0,T], L_1)$. Assertion (2.29) is now evident. \square

Remark 2.9. *Theorem 2.8 implies that there exist no further equilibria in L_1^+ for which V is finite.*

Remark 2.10. *Note that it holds $V(w) < \infty$ for $w \in L_p^+$ provided $p > 1$. This is a consequence of the properties of H , Hölder's inequality, and the fact that*

$$x|\log x| \leq c(\varepsilon)(x^{1+\varepsilon} + x^{1-\varepsilon}) , \quad x > 0 , \quad \varepsilon > 0 .$$

Remark 2.11. Observe that the asymptotic distribution provided by Theorem 2.8 depends merely on the total mass of the initial distribution but not on its shape, which seems to be consistent with numerical simulations and physical theory (see [14], [15], [21], and [25] for details).

It may be worthwhile to present some examples of kernels satisfying the imposed assumptions.

Example 2.12. If φ is defined as in (1.5), then hypothesis (H_1) is satisfied provided $\Phi : \mathbb{R}^2 \rightarrow (0, \infty)$ is uniformly Lipschitz continuous on bounded sets and bounded.

Example 2.13. Let $P \in C((0, y_0]^2, (0, \infty))$ be symmetric and let $q \in C((0, y_0], \mathbb{R}^+)$ be such that

$$0 < P(y, y') + q(y + y') \leq 1, \quad 0 < y + y' \leq y_0.$$

Assume $\alpha \geq 0$ and $0 \geq \alpha - \beta > -1$ and define for arbitrary constants $K^*, \gamma^* > 0$

$$\begin{aligned} Q(y, y') &:= q(y + y'), \quad 0 < y + y' \leq y_0, \\ K(y, y') &:= K^*(y + y')^\alpha, \quad 0 < y, y' \leq y_0, \\ \gamma(y, y') &:= \gamma^* P(y - y', y') y^\beta [y'(y - y')]^{\alpha - \beta}, \quad 0 < y' < y \leq y_0, \\ \beta_c(y, y') &:= c_{\alpha, \beta} y^{-1 - 2\alpha + 2\beta} [y'(y - y')]^{\alpha - \beta}, \quad 0 < y' < y \leq y_0, \\ \beta_s(y, y') &:= f_s(y) [y'(y - y')]^{\alpha - \beta}, \quad 0 < y - y_0 \leq y' \leq y_0, \end{aligned}$$

where $c_{\alpha, \beta} := (\mathbf{B}(\alpha - \beta + 2, \alpha - \beta + 1))^{-1}$ with \mathbf{B} denoting the beta function and where

$$f_s(y) := y \left(\int_{y - y_0}^{y_0} y' [y'(y - y')]^{\alpha - \beta} dy' \right)^{-1}, \quad y_0 < y < 2y_0.$$

Then hypotheses $(H_2) - (H_6)$ are satisfied with

$$H(y) := \frac{\gamma^*}{K^*} y^{\alpha - \beta}, \quad y \in (0, y_0].$$

Further, (2.28) holds provided $\alpha = \beta$.

Example 2.14. Analogously as in [16] we may define

$$\begin{aligned} K(y, y') &:= r e^{-y^2 - (y')^2}, \quad 0 < y, y' \leq y_0, \\ \gamma(y, y') &:= s e^{-(y - 2y')^2}, \quad 0 < y' < y \leq y_0, \\ \beta_s(y, y') &:= f(y) e^{-4y(y - y')}, \quad 0 < y - y_0 \leq y' \leq y_0, \end{aligned}$$

for some $r, s > 0$, where

$$f(y) := y \left(\int_{y - y_0}^{y_0} y'' e^{-4y(y - y'')} dy'' \right)^{-1}, \quad y_0 < y < 2y_0.$$

Then, for $P \equiv 1$ and $Q \equiv 0$, hypotheses $(H_2) - (H_6)$ hold with

$$H(y) := \frac{s}{r} e^{-y^2}, \quad y \in (0, y_0],$$

and, in addition, (2.28) is satisfied.

Example 2.15. The other example from [16] can also be considered. Let α, τ, p , and λ be arbitrary real numbers and let $A_0, B_0 > 0$. Put

$$\begin{aligned} K(y, y') &:= A_0(1+y)^\alpha(1+y')^\alpha, \\ \gamma(y, y') &:= B_0K(y', y-y')(1+y)^\tau[(1+y')(1+y-y')]^{-\tau}e^{\lambda(y^p-(y-y')^p-(y')^p)}, \\ \beta_s(y, y') &:= y\nu(y, y')\left(\int_{y-y_0}^{y_0} y''\nu(y, y'') dy''\right)^{-1}, \end{aligned}$$

where $\nu(y, z) := (1+z)^{\alpha-\tau}(1+y-z)^{\alpha-\tau}e^{-\lambda(z^p+(y-z)^p)}$. Then, with $P \equiv 1$, $Q \equiv 0$, and

$$H(y) := \frac{B_0}{A_0}(1+y)^{-\tau}e^{-\lambda y^p-y}, \quad y \in (0, y_0],$$

hypotheses $(H_2) - (H_6)$ and inequality (2.28) are satisfied.

3. Stability

We now focus on stability of the equilibria. For this purpose let us introduce for any $\varrho > 0$ the spaces

$$X^+ := \{u \in L_1^+; V(u) < \infty\} \quad \text{and} \quad X_\varrho^+ := \{w \in X^+; M(w) = \varrho\}.$$

If not stated otherwise, X^+ and X_ϱ^+ are equipped with the L_1 -topology turning them into metric spaces. Observe that both X^+ and X_ϱ^+ are positively invariant, and that the map $(t, u^0) \mapsto u(t; u^0)$ defines a semiflow on X^+ and X_ϱ^+ due to Theorem 2.1 and Proposition 2.7. Moreover, provided (2.28) holds, Theorem 2.8 entails that $u_{\alpha(\varrho)}$ is a global attractor for the semiflow generated on X_ϱ^+ , where $\alpha(\varrho)$ is chosen such that $M(u_{\alpha(\varrho)}) = \varrho$.

In order to state the next proposition, we define for $\eta \in \mathbb{R}$

$$V_\eta(w) := V(w) - |H|_1 - \eta M(w), \quad w \in X^+.$$

Proposition 3.1. For $\varrho > 0$ choose $\alpha(\varrho) \in \mathbb{R}$ such that $M(u_{\alpha(\varrho)}) = \varrho$. Then, $u_{\alpha(\varrho)}$ is the unique minimizer of V on X_ϱ^+ and of $V_{\alpha(\varrho)}$ on X^+ . Moreover, for any minimizing sequence (w_j) of V on X_ϱ^+ , it holds $w_j \rightarrow u_{\alpha(\varrho)}$ in X_ϱ^+ .

Proof. For $r > 0$ define

$$f_r(w) := w\left(\log \frac{w}{r} - 1\right), \quad w \geq 0,$$

with $f_r(0) := 0$. Then f_r has at $w = r$ a global minimum for each $r > 0$. For brevity put $\alpha := \alpha(\varrho)$. Given $w \in X^+$ it holds

$$V_\alpha(w) = \int_0^{y_0} f_{u_\alpha(y)}(w(y)) dy \geq \int_0^{y_0} f_{u_\alpha(y)}(u_\alpha(y)) dy = V_\alpha(u_\alpha),$$

where the inequality is strict if w differs from u_α on a set of non-zero measure. Hence, u_α is the unique minimizer of V_α on X^+ . Furthermore, since $M(X_\varrho^+) = \{\varrho\}$, it also

minimizes V on X_ϱ^+ .

Let now (w_j) be a minimizing sequence of V in X_ϱ^+ , i.e.

$$\lim V(w_j) = \inf_{w \in X_\varrho^+} V(w) = V(u_\alpha) . \quad (3.1)$$

Observing that this implies

$$|f_{u_\alpha(\cdot)}(w_j(\cdot)) - f_{u_\alpha(\cdot)}(u_\alpha(\cdot))|_1 = V_\alpha(w_j) - V_\alpha(u_\alpha) \longrightarrow 0 ,$$

we may extract a subsequence (j') such that $f_{u_\alpha(\cdot)}(w_{j'}(\cdot)) \rightarrow f_{u_\alpha(\cdot)}(u_\alpha(\cdot))$ a.e.. This easily implies $w_{j'} \rightarrow u_\alpha$ a.e.. From (3.1), Lemma 2.2, and the Dunford-Pettis theorem we deduce that $(w_{j'})$ is relatively weakly compact in L_1 . Therefore, there exists a further subsequence (j'') and $w \in L_1$ such that $w_{j''} \rightarrow w$ in $L_{1,w}$. Since V is weakly lower semi-continuous we get

$$V(w) \leq \liminf_{j''} V(w_{j''}) = V(u_\alpha) < \infty ,$$

whence $w \in X_\varrho^+$. From the above considerations we conclude $w = u_\alpha$. Altogether, we obtain $w_{j''} \rightarrow u_\alpha$ in $L_{1,w}$ and a.e. so that $w_{j''} \rightarrow u_\alpha$, from which the assertion follows. \square

Theorem 3.2. *Let $\varrho > 0$ be given and choose $\alpha(\varrho) \in \mathbb{R}$ such that $M(u_{\alpha(\varrho)}) = \varrho$. Then, for each $\varepsilon > 0$ there exists $\delta > 0$ such that for any $u^0 \in X_\varrho^+$ with*

$$|u^0 - u_{\alpha(\varrho)}|_1 < \delta \quad \text{and} \quad V(u^0) < V(u_{\alpha(\varrho)}) + \delta$$

it holds $|u(t; u^0) - u_{\alpha(\varrho)}|_1 < \varepsilon$ for $t \geq 0$.

Proof. Due to [3, Prop.4.3] we merely have to show that V is decreasing along orbits — which was done in Proposition 2.7 — and that $u_{\alpha(\varrho)}$ lies in a 'potential well' with respect to X_ϱ^+ , that is, for given small $\varepsilon > 0$ there exists $\sigma(\varepsilon) > 0$ such that $V(w) - V(u_{\alpha(\varrho)}) \geq \sigma(\varepsilon)$ for all $w \in X_\varrho^+$ with $|w - u_{\alpha(\varrho)}|_1 = \varepsilon$. But this readily follows from Proposition 3.1. \square

Define the metric d by

$$d(w, v) := |w - v|_1 + |V(w) - V(v)| , \quad w, v \in X^+ .$$

We conclude with a stability result being a stright consequence of the decrease of V along orbits.

Corollary 3.3. *Let $\varrho > 0$ be arbitrary and choose $\alpha(\varrho) \in \mathbb{R}$ such that $M(u_{\alpha(\varrho)}) = \varrho$. Then, the equilibrium $u_{\alpha(\varrho)}$ is stable in (X_ϱ^+, d) , that is, for each $\varepsilon > 0$ there exists $\delta > 0$ such that for any $u^0 \in X_\varrho^+$ with $d(u^0, u_{\alpha(\varrho)}) < \delta$ it holds $d(u(t; u^0), u_{\alpha(\varrho)}) < \varepsilon$ for $t \geq 0$.*

Remark 3.4. *For the case without scattering and shattering it is shown in the recent paper [17] that*

$$V(u(t; u^0)) \rightarrow V(u_\alpha) \quad \text{as } t \rightarrow \infty ,$$

where $M(u_\alpha) = M(u^0)$. Such an improvement of Theorem 2.8 would allow to conclude asymptotical stability of the equilibrium $u_{\alpha(\varrho)}$ in (X_ϱ^+, d) .

Acknowledgements

This research consists of an edited extract of the author's Ph.D. thesis submitted to the Universität Zürich. The author gratefully acknowledges the support of his thesis supervisor Prof. H. Amann.

References

- 1 M. Aizenman and T.A. Bak. Convergence to equilibrium in a system of reacting polymers. *Comm. Math. Phys.* **65** (1979), 203-230
- 2 J.M. Ball, J. Carr and O. Penrose. The Becker-Döring cluster equations: Basic properties and asymptotic behaviour of solutions. *Comm. Math. Phys.* **104** (1986), 657-692
- 3 J.M. Ball and J.E. Marsden. Quasiconvexity at the boundary, positivity of the second variation and elastic stability. *Arch. Rat. Mech. Anal.* **86** (1984), 251-277
- 4 I. Borsi. Dynamics of liquid-liquid dispersions with unbounded fragmentation kernel. *Adv. Math. Sci. Appl.* **11**, No. 2 (2001), 571-591
- 5 J. Carr. Asymptotic behaviour of solutions to the coagulation-fragmentation equations. I. The strong fragmentation case. *Proc. R. Soc. Edinb.* **121A** (1992), 231-244
- 6 J. Carr and F.P. da Costa. Asymptotic behaviour of solutions to the coagulation-fragmentation equations. II. Weak fragmentation. *J. Stat. Phys.* **77**, No. 1/2 (1994), 89-123
- 7 Z. Cheng and S. Redner. Scaling theory of fragmentation. *Phys. Rev. Lett.* **60**, No. 24 (1988), 2450-2453
- 8 Z. Cheng and S. Redner. Kinetics of fragmentation. *J. Phys. A: Math. Gen.* **23** (1990), 1233-1258
- 9 J.F. Collet and F. Poupaud. Asymptotic behaviour of solutions to the diffusive fragmentation-coagulation system. *Physica D* **114** (1998), 123-146
- 10 F.P. da Costa. Convergence to equilibrium of solutions to the coagulation-fragmentation equations. In *Nonlinear Evolution Equations and their Applications* (Macau, 1998), 45-56, World Sci. Publishing, River Edge, NJ 1999
- 11 R.E. Edwards. *Functional Analysis. Theory and Applications.* (Dover Publ. 1995)
- 12 I. Ekeland and R. Temam. *Analyse convexe et problèmes variationnels.* (Dunod 1974)
- 13 A. Fasano and F. Rosso. A new model for the dynamics of dispersions in a batch reactor: theory and numerical simulation. In *Lectures on applied mathematics: proceedings of the symposium on the occasion of Karl-Heinz Hoffmann's 60th birthday, Munich, June 30 - July 1, 1999.* (ed. H.J. Bungartz, R. Hoppe and C. Zeuger), pp. 123-141 (Springer 2000)
- 14 A. Fasano. The dynamics of two-phase liquid dispersions: necessity of a new approach. *Milan J. Math.* **70** (2002), 245-264
- 15 M. Kostoglou and A.J. Karabelas. An explicit relationship between steady-state size distribution and breakage kernel for limited breakage processes. *J. Phys. A: Math. Gen.* **30**, No. 20 (1997), L685-L691
- 16 P. Laurençot and S. Mischler. The continuous coagulation-fragmentation equations with diffusion. *Arch. Rat. Mech. Anal.* **162**, No. 1 (2002), 45-99
- 17 P. Laurençot and S. Mischler. Convergence to equilibrium for the continuous coagulation-fragmentation equation. *Bull. Sci. math.* **127** (2003), 179-190
- 18 P. Laurençot and D. Wrzosek. Fragmentation-diffusion model. Existence of solutions and their asymptotic behaviour. *Proc. R. Soc. Edinb.* **128** (1998), 759-777
- 19 P. Laurençot and D. Wrzosek. The Becker-Döring model with diffusion: II. Long time behaviour. *J. Diff. Equ.* **148** (1998), 268-291
- 20 P. Laurençot and D. Wrzosek. The discrete coagulation equations with collisional breakage. *J. Statist. Phys.* **104**, No. 1-2 (2001), 193-253
- 21 A. Mancini and F. Rosso. A new model for the dynamics of dispersions in a batch reactor: numerical approach. *Meccanica* **37**, No. 3 (2002), 221-237
- 22 K. Panoussopoulos. Separation of crude oil-water emulsions: experimental techniques and models. Ph.D. Thesis. ETH Zürich (1998)
- 23 I.W. Stewart and P.B. Dubovskii. Approach to equilibrium for the coagulation-fragmentation equation via a Lyapunov functional. *Math. Meth. Appl. Sci.* **19** (1996), 171-183

- 24 I.W. Stewart and P.B. Dubovskii. Trend to equilibrium for the coagulation-fragmentation equation. *Math. Meth. Appl. Sci.* **19** (1996), 761-772
- 25 K. Valentas, O. Bilous and N.R. Amundson. Breakage and coalescence in dispersed phase systems. *I&E C Fundamentals* **5** (1966), 533-542
- 26 I.I. Vrabie. *Compactness methods for nonlinear evolutions*. Second edition. (Longman 1995)
- 27 C. Walker. Coalescence and breakage processes. *Math. Meth. Appl. Sci.* **25** (2002), 729-748
- 28 C. Walker. On diffusive and non-diffusive coalescence and breakage processes. Ph.D. Thesis. Universität Zürich (2003)
- 29 D. Wilkins. A geometrical interpretation of the coagulation equation. *J. Phys. A: Math. Gen.* **15** (1982), 1175-1178